Newborn Screening

Oklahoma State Department of Health
Newborn Screening Program

Phone: (405) 426-8310

Toll Free: 1 (800) 766-2223

Fax: (405) 900-7556

NewbornScreen@health.ok.gov



Why Newborn Screening

Allow me to Introduce....





Jase's Story – Galactosemia



History of Newborn Screening


Then and Now






History of Newborn Screening

- Newborn screening originated with Dr. Robert Guthrie who developed a test for elevated phenylalanine in dried blood spots in 1960. (PKU Disease)
- Before the blood test existed, most children with PKU were not diagnosed until after they had irreversible brain damage.
- Early test > Early diagnosis > Early treatment > Mitigated brain damage.
- PKU was the first condition identified by NBS, so some people still refer to all NBS as the "PKU test." However, this term is not accurate as the newborn screen now tests for 57 total disorders, not solely PKU.






Before NBS: Parents Had to Lose One to Save One



Untreated versus Treated PKU

Scientific Progress Translates to More Infant Lives Positively Impacted

Newborn Screening Timeline





Newborn Screening Today: A Three Part Process

- Newborn screening checks a baby for certain conditions present at birth that benefit from early treatment or intervention.
 - Blood spot screening, which determines if a baby might have one of many serious conditions
 - Pulse oximetry screening, which determines if a newborn might have certain heart conditions
 - Hearing screening, which determines if a newborn might be deaf or hard of hearing











PURPOSE – Every Baby Deserves A Newborn Screen

- Newborn screening (NBS) is the practice of testing **every** newborn for harmful or potentially fatal disorders that are not otherwise apparent at birth.
- Early detection and prompt treatment can make the difference between healthy development or lifelong impairment and possible death.





Screening VS Diagnostic

• Screening results, by themselves, cannot determine the presence or absence of a disorder. The purpose of a screen is to detect risk factors for disease in large numbers of apparently healthy individuals – in this case, newborns.


• **Diagnostic results** refers to the combination of signs, symptoms, and test results that allows the provider to confirm the diagnosis of the respective disease.





Who Decides?


- In 2022 the Oklahoma legislature passed statute stating that the Oklahoma NBS panel will match the national <u>Recommended Uniform Screening Panel</u> (RUSP) to the extent practicable.
- Once a condition is added to the RUSP the NBS Program (lab and follow up) will determine practicability and readiness.
- The Infant and Children's Health Advisory Committee will provide recommendations to the Commissioner of Health to add the disorder.
- The Commissioner of Health will give final approval.
- Oklahoma currently screens for 57 possible hidden disorders.
- Oklahoma will continue to expand.
- 3 disorders added to RUSP in the last year (MPS2, GAMT, and Krabbe) that OK will work to add to our state panel.



Autosomal Recessive

- Most NBS disorders are autosomal recessive with the exception of:
 - Congenital Hypothyroidism (CH)
 - Some forms of Severe Combined Immunodeficiency (SCID)
 - X-Linked Adrenoleukodystrophy
- Usually no prior family history
- Risk for <u>each</u> pregnancy if both parents are a carrier of a disorder:


Possible Outcomes for Offspring of Parental Disease Carriers






Parent Education

- Instruct parents to ask for their baby's newborn screening results:
 - Baby's Pediatrician
 - Local County Health Department
 - OSDH Newborn Screening Program
- Tell parents to hang onto the Blue or Pink slip from their baby's filter paper for reference.



Oklahoma State Rules and Statutes – Filter Paper Education Pages Blood spot



Every baby born in Oklahoma is required to have blood tests performed during the first week of life in order to help in the early detection of a group of treatable medical conditions that can cause severe illness, developmental disability or death. These tests can all be performed using a small amount of blood usually collected when the baby is 24 to 48 hours old. The blood sample is sent to the Oklahoma State Department of Health (OSDH) Public Health Laboratory for testing. Test results are usually available in 10-14 days. For a list of conditions that are screened for in Oklahoma, see the OSDH Newborn Screening Program website at http://nsp.health.ok.gov

Importance of newborn screening

A baby with one of the conditions in the newborn screening test panel may appear healthy at birth, which makes it difficult for health-care providers to recognize clinically. Failure or delay in diagnosing and treating a baby with one of these conditions within weeks of life can lead to severe illness or death. Newborn screening blood tests help inform healthcare providers if your baby is at risk for one of these conditions. If your baby is found to have a disorder, immediate care by a medical specialist may be needed.

How will I get the test results for my baby?

Please, take this form with you to your baby's first well child visit and ask for your baby's newborn screening test results. If your baby's healthcare provider does not have the test results and you have not been notified by mail, please call the OSDH Newborn Screening Program at the number indicated on the reverse of this form when your baby is 3 weeks of age.

OKLAHOMA NEWBORN SCREENING PROGRAM

Oklahoma State Department of Health

Parent/Guardian Information Sheet

Will my baby need more testing?

Your baby's healthcare provider or an OSDH Newborn Screening Program coordinator will contact you if your baby needs further testing. They will tell you why more tests are needed and what to do next. Retesting does not necessarily mean that your baby is sick, but rather is done to be sure there is not a problem.


Additional testing may be needed if:

- Test results were abnormal or unclear.
- Your baby was premature or sick at birth.
- The blood sample was collected before your baby was 24 hours of age.
- Your baby had a blood transfusion before the blood sample was collected.
- There was a problem with the blood sample.
- · Your baby's healthcare provider requests repeat testing.

What if I have questions?

If you have questions about your baby's newborn screening tests or test results, contact your baby's healthcare provider, visit the OSDH Newborn Screening Program website at http://nsp.health.ok.gov, call the OSDH Newborn Screening Program at (405) 271-6617 or 1-800-766-2223 or email the program at newbornscreen@health.ok.gov

Oklahoma State Rules and Statutes – Filter Paper Education Pages Hearing Screen



OKLAHOMA NEWBORN HEARING SCREENING PROGRAM

Oklahoma State Department of Health


Parent/Guardian Information Sheet

Your baby's hearing

Your child's most important learning and speech development will take place during the first few years of life. In these early years of development, your child learns how to communicate — first to understand what people say, and then to start talking. Any degree of undetected hearing loss can negatively impact a child's speech, language, social and emotional development.

Your baby should be able to achieve the following milestones around the ages listed below. As the weeks and months go by, check to see if your baby can do most of the things listed. If your baby can't, don't wait— have your infants' hearing tested. If you suspect a hearing loss or have a concern about your child's hearing, contact your healthcare provider, an audiologist, or your county health department to find out about hearing testing.

Hearing checklist



What if I have questions?

If you have questions about your baby's newborn hearing test results, contact your baby's healthcare provider, visit the OSDH Newborn Screening Program website at http://nsp.health.ok.gov, call at (405) 271-6617 or 1-800-766-2223, or email the program at newbornscreen@health.ok.gov.

Parent Education

- NBS is collected on **every** baby born in Oklahoma.
- Importance of correct contact info & PCP for follow-up.
- No news is not good news! Update NBS Program with changes in home address and/or PCP.
- Review hidden disorders, using NBS pamphlet as a guide.
- Specimens are kept by the OSDH lab for 42 days before being destroyed.
- Explain that most affected newborns do not exhibit signs & symptoms early on.
- Prompt identification & treatment of disorders is critical.

Indications for Repeat Screen

- The NBS testing results are out of range for one or more disorders PCP notified by phone, parent by mail.
- The NBS sample was damaged or is otherwise unsatisfactory for testing PCP and parent notified by mail.
- The infant received a transfusion prior to NBS collection usually in NICU, will notify PCP when time for recollection.
- The infant's screen was collected prior to 24 hours old if normal result, this report will be on the portal, if out of range, PCP notified by phone, parent by mail.
- The infant is premature or sick (TPN and antibiotics could affect results.




Filling Out the Demographic Form






Filling out the Form






Filling out the Form

Specimen testing will be delayed if the form is incomplete!




Check expiration date

- If the filter paper is expired, discard the paper, and check the stock of filter paper kits and discard all expired kits.
- Collect the specimen on a kit that is not expired.



Specimen testing will be delayed if the form is incomplete!




If you don't mark "repeat screen" the correct linking to the original screen could be delayed.

- If this is the first specimen collected for the baby, check the "First Screen" box.
- If baby has had a previous screen, check the "Repeat Screen" box.
 - List the previous OSDH Lab Number, if you have it.








- If baby expires before a screen can be collected:
 - Check the "Expired" box
 - Enter the date that baby passed away
 - Submit the filter paper form to the OSDH PHL





Specimen testing will be delayed if the form is incomplete!




If baby is transferred prior to specimen collection:

- Check the "Transferred" box
- Enter the date that baby transferred and the facility that baby was transferred to
- It is the responsibility of the receiving facility to collect the newborn screen





Specimen testing will be delayed if the form is incomplete!



BABY'S INFORMATION

- Tests Requested: Check all that apply.
 - All Tests- always check unless test is for HGB
 Only. This ensures the lab screens for all
 disorders on the NBS panel.
 - HGB Only- Check if repeat screen is for follow-up of initial abnormal HGB result.
 - GALT- Check GALT in addition to All Tests if there
 is a family history of galactosemia or if baby is on
 lactose-free (soy) formula at time screen is
 collected.
 - Phe Monitor- Check only if baby has been diagnosed with PKU (typically metabolic specialists only)
 - **CFTR** Check in addition to All Tests if baby has clinical concerns for CF, meconium ileus, and/or family history of CF.

Transferred

Filling out the Form: Infant's Information – Initial screen




- Baby's first and last name (use legal name as displayed on the birth certificate).
- If baby's first name is unknown, "BG" or "Female", "BB" or "Male" may be used.


	BACY'S INFORM	TION			
Last Name	First Name				
Birth Date//	Time:(24 Hr Clock) Male	Race (Check all that apply) White		
Collection Date//		Female Unknown	Black Hispanic Asian		
Medical Record #	Gest. Age Birth Wt. (gm)	Multiple Birth Order A-H	American Indian Pacific Islander		



Filling out the Form: Infant's Information – for repeat screen




- Baby's first and last name
- If baby's name was updated after hospital discharge, screens will be linked using the DOB, previous specimen #, mom's name, address and/or phone.









- Sex/Gender
 - Check "Male", "Female", or "Unknown"





- Date & <u>Time</u> of birth
 - Enter the time using the 24 hour clock. For example 1PM would be entered as 13:00.
 - For a repeat screen, outside of hospital, if time is not known, this box can be left empty.







- <u>Date</u> & <u>Time</u> of specimen collection
 - Ideal time for well, term newborn:

24 hours + 1minute of age


- Enter the time using the 24 hour clock. For example 1PM would be entered as 13:00.
- Ideal date of repeat is determined by follow up recommendations.








- Medical record number
 - Baby's medical record number
 - If a multiple birth, take extreme care here




Filling out the Form: Infant's Information



- Gestational Age
 - List gestational age at birth, may leave blank on a repeat collection.
 - Lab cut off values for abnormal severe combined immunodeficiency (SCID) are gestational age dependent.



Filling out the Form: Infant's Information



- Birthweight (in grams), leave blank if unknown (for repeat screen out of hospital)
 - Lab cut off values for abnormal congenital adrenal hyperplasia (CAH) results are dependent on birth weight.



Filling out the Form: Infant's Information



- Birth order (if multiple birth is present)
 - Indicate "A", "B", "C", etc. if baby is of multiple birth (twin, triplet, etc.).
 - Do <u>NOT</u> mark anything in this space if baby is a single birth.

BABY'S INFORMATION							
Last Name		First Name					
		S	ex	Race (Check all that apply)			
Birth Date / /	Time:	(04.11-01-11	7	White			
birti Date/	nme	(24 Hr Clock)	Male				
		L	Female	Black			
0-115 0-1 1 1			Unknown	Hispanic			
Collection Date//	Time	(24 Hr Clock)		Asian			
Medical Record #	Gest. Age Birth Wt.	(gm)		American Indian			
	1 1	Multiple	A-H				
		Birth Order		Pacific Islander			



Filling out the Form: Mom's Information

Specimen testing will be delayed if the form is incomplete!



Note: If baby is <u>adopted</u>, be sure to check the <u>Adoption</u> box on the filter paper form and enter the agency/law firm information in this section. If <u>DHS</u> is involved, enter case worker information in this section and check the <u>DHS</u> <u>Custody</u> box.

• DHS Custody or Adoption



Filling out the Form: Mom's Information



- Mom's first and last name
- Mom's mailing address:
 - Street, Apt # (if applicable), City, State, Zip
- Mom's telephone number:
 - Extremely important to include in case newborn screen results are abnormal and require followup.






- Physician Ordering the NBS:
 - Include first and last name
 - Enter the NBS Provider ID #, if known
 - May leave blank if ordering Physician unknown on a repeat collection, outside of hospital




Filling out the Form: Provider's Information Specimen testing will be delayed if the form is incomplete!




- Primary Care/Follow-up Physician:
 - Planned health care provider upon discharge from birthing facility
 - Include first and last name
 - Enter the NBS Provider ID #, if known
 - <u>Extremely important</u> that this is correct in case newborn screen results are abnormal and require follow up



Filling out the Form: Medical/Feeding History Specimen testing will be delayed if the form is incomplete!





Check all that apply for baby at the time of specimen collection

- If transfused enter the <u>date</u> and <u>time</u> of transfusion
- NICU/Special Care Nursery
- TPN/SNAP
- Lipids/Carnitine/MCT
- Lactose-Free (Soy) Formula




- Meconium Ileus
- Family History of Cystic Fibrosis (CF)




Filling out the Form: Submitter ID


Specimen testing will be delayed if the form is incomplete!



- Submitting Health Provider ID #
 - This is the ID of the provider/facility who collected the specimen
 - Write or stamp in facility name and address



Filling out the Form: Collector's Initials Allows for thorough follow up of an unsatisfactory screen



Note: Do not touch the filter paper in any other area when writing initials and unit.

Unsatisfactory Specimen Follow-up


 Specimen collectors can place their initials and unit in the area below for identification purposes, in the event of an unsatisfactory specimen. This allows for easier identification of the individual who collected the specimen so that further education and/or training can be provided.

Collecting the Specimen






How Is An Infant Screened?



- Blood spot screen-heel stick
- Performed when the infant is "24 hours plus one minute" or prior to discharge, whichever comes first.
- Small amount of blood is placed on a small card and sent to the OSDH Newborn Screen lab
- Newborn screening specimens are picked from birthing hospitals and county health department via a contract courier service and then brought to the PHL for testing.






Time of Screening: Healthy Newborn

"24 hours plus one minute" of age
Or
Prior to discharge

WHICHEVER COMES FIRST





Time of Screening: Premature or Sick Newborns



Specimen Collection


- Heel Stick / Direct Application
 - Preferred, recommended method
 - Start with clean, dry hands before handling the filter paper.






Prepare the Site

- Warm the heel with a heel warmer or a soft cloth, moistened with warm water up to 41 C for 3-5 min.
- Warmth leads to vasodilation, which increases blood flow and chance of collection success.
- Follow your facility protocol regarding which warming device to use.




Prepare the Site

- If desired, parent may hold infant during collection
 - Decreases stress response in newborn
 - Encourages bonding
- Position the infant's leg lower than the heart.
 - This increases venous pressure, which results in increased blood flow and a greater chance of collection success.
- Wearing gloves, wipe the infant's heel with 70% isopropyl alcohol.
- Allow the heel to air dry!
 - Residual alcohol can affect NBS results and/or lead to unsatisfactory specimens.




Lancet Placement

- Hatched areas are safe for puncture
 - Damage to nerves and/or the heel bone may occur for punctures outside of the hatched region.



Perform the Puncture

- Using a sterile lancet, perform the puncture.
- Gently wipe off the first drop of blood with a sterile gauze or cotton ball.
- Apply gentle pressure with thumb and around heel but not near the puncture site; ease intermittently as drops of blood form.
- Avoid "milking" the puncture site.




Application

Gently touch the filter paper card to the blood drop and fill each printed circle with **one** large drop of blood.

- Apply blood to one side only.
- Observe the saturation of each printed circle as the blood flows through the filter paper.


What about capillary tubes? >> Not preferred

- Higher risk for collection error
- If used, must be sterile/clean and plain.
- No additives, must be anticoagulant free, however no anti-coagulants = risk of clotting.
- Risk of scratching the filter paper, avoid touching tip of tube to the paper.
- Use a new tube for EACH printed circle.



What about venous samples? >> Discouraged

- May be appropriate under certain circumstances (e.g. NICU)
- More invasive than a heel stick.
- Do not draw blood from extremity with infusing IV fluids.
- Please refer to CLSI guidelines for more information.



Only in certain circumstances

What about umbilical catheters? >> Discouraged

- May be appropriate under certain circumstances (e.g. NICU).
- Ensure the line is cleared with withdrawing 2-2.5 cc (ml) of blood to collecting a specimen for newborn screen.
- Please refer to CLSI guidelines for more information.



What about umbilical cord blood? >> Discouraged

- May be appropriate under certain circumstances (e.g. NICU)
- Risk for maternal blood contamination.
- Repeat the newborn screen using the heel stick method when indicated.
- Please refer to CLSI guidelines for more information.

Specimen Collection: What NOT to Do

- Do NOT dab or "color in" the filter paper circles.
- Do NOT apply multiple drops of blood per circle.
- Do NOT scratch the filter paper.
- Do NOT contaminate specimens.
 - insufficient drying of alcohol, oils on hands, lotions, compressing the circles, spills, etc..
- Do NOT stack specimens.
 - risk for leaching & cross-contamination between specimens
- Do NOT submit wet specimens.
- Do NOT place specimens in direct sunlight or in front of air vents or other sources of moving air.
- Do NOT place specimens in plastic bags.
- Do NOT batch (hold onto) specimens.



Collection Reminders

Pre-collection:

not expired.


• Check the Expiration Date of the filter paper

If filter paper is expired, discard the paper, check the stock of filter paper kits it came from to ensure they are not all expired, and collect on a kit that is

Post-collection:


- Air dry specimen horizontally for 3-4 hours
 - Transporting wet specimens can make them unsatisfactory for testing.
- Send specimen with Courier within 24 hours of collection
 - Delayed receipt of specimens to the Public Health Laboratory can delay identification of and treatment for a disorder, which can result in lifelong disability or even death for Oklahoma newborns.
 - Know the courier schedule and location for your facility! Ensure all staff involved in newborn screening are also aware of the process.
- Maintain specimen collection log & ensure screening results are received & recorded
- Ensure that everybody who handles the filter paper or is involved in the newborn bloodspot collection process is trained

A Peek Inside the Laboratory.....











Inside the Laboratory







Blood Spot Screening is a system, not an event.



Diagnostic
Testing,
Potential
Diagnosis and
Long Term
Follow-Up

^{*}Short-term Follow-up continues tracking case until confirmed normal or diagnosed with a disorder.

NBS Filter Paper Review


Unsatisfactory (Unsat) Specimen Examples





Filter Paper

- The filter paper is part of the NBS Form. It is a medical device designed to absorb a specific volume of blood within each pre-printed filter paper circle.
- If an analyte for any disorder is either too high or too low, this is an indication that additional testing is needed.
- Accurate results depend upon proper absorption of blood onto the filter paper.
 - Too much or too little blood may result in inaccurate results.




Multiple Application



Why Unsat?

- When bloodspots overlap or touch, as is the case in the sample above, it creates an uneven absorption of blood.
- Analyte levels cannot be accurately measured.
- Testing these specimens will result in inaccurate results.


Clotted or Caked Blood



Why Unsat?


- Clots can occur using capillary tubes or if too much blood is applied to the pre-printed circles.
- Samples with clots are not suitable for testing.

Serum Rings




Why Unsat?

- Notice the halos around the periphery of most of the pre-printed circles above. This can occur due to the following:
 - Insufficient drying of alcohol on the baby's heel prior to heelstick
 - Drying the specimen vertically instead of horizontally
 - Closing the flap of the filter paper on top of the circles while the specimen is still wet
 - Placing wet specimens in plastic bags
 - Milking or squeezing the puncture site




Inadequate Amount of Blood



Why Unsat?

• The above filter paper circles are not sufficiently filled with blood for testing.


Under-Saturation



Why Unsat?

• Notice how the blood has not soaked all the way through the filter paper. There simply is not enough blood in this sample for testing.

Acceptable Filter Paper



Why Acceptable?

- Pre-printed circles are completely filled with blood
- Blood has soaked all the way through the filter paper
- Absence of clots or caked blood
- Absence of serum rings



Are All 5 Circles Needed?





Why?

- If a result is flagging out-of-range, the specimen will be retested and the final result with be an average of three results. Each test requires an additional punch to be taken from the pre-printed circles.
- If the results for Congenital Adrenal Hyperplasia (CAH), Pompe Disease, MPS1 or X-ALD are out-of-range, *one or two entire circles* will be removed & shipped to another laboratory for steroid profile testing. These are overnighted to Mayo laboratory in Minnesota or Revvity Labs in Massachusetts. Screen results are not final until second tier is complete
- Disorders will continue to be added to the newborn screening panel.
- The specialist and family may request for the specimen to be sent to another laboratory for additional testing to assist in determining diagnosis.

For Reference...

• Refer to *Clinical and Laboratory Standards Institute* (CLSI) for collection guidelines.



NICU and Special Considerations





NICU Special Considerations



- Prematurity & LBW may affect TSH & 17-OHP results
- Hypoxia, CMV, septicemia, trisomies, biliary atresia may affect IRT levels
- Liver immaturity may affect amino acid results
- Carrier status may affect all NBS results



- TPN, SNAP, & carnitine may affect amino acid, fatty acid, or organic acid results
- Steroids may affect 17-OHP results
- ECLS & blood transfusions may affect all NBS results



- PTU therapy or radioactive iodine may affect infant TSH results
- Steroids may affect infant 17-OHP results



- Contamination: oils/lotion from hands, spills, standing water, residual alcohol, heat/humidity
- Early/delayed specimen collection
- Transit time delays
- Unsatisfactory specimens

Additional Information






Hospital Responsibilities

- Ensure **ALL** infants are screened prior to discharge. (Keep a log book to ensure compliance.)
- Ensure specimens are received in a timely manner to the OSDH PHL for testing.
- Infants who are transferred: >> The **RECEIVING** hospital is to ensure that the NBS has been collected.
- Submit **SATISFACTORY** specimens:
 - Collected properly
 - All requested information is documented on the demographic form attached to filter paper
 - Submitted timely



Data used for hospital reporting is pulled by the date the OSDH laboratory received the specimen. Specimen types included in hospital reporting are: Less Than 24 Hour Specimens, Initial Specimens, & Repeat Specimens.






Refusal of screening

- Parents may refuse screens based on Religious Tenets and Practices.
- Refusal form must be signed and placed in the medical record with a copy mailed to the Newborn Screening Program.
- Please note that parent must indicate which of 3 portions of screen are being refused: Hearing, bloodspot, and/or CCHD.
- Demographic form with no specimen on filter paper should be sent into PHL with courier, with "X" marked in refusal box.



Refusal

- Religious Tenets and Practices only.
- Check the box(es) on the filter paper form if parents refuse the NBS and/or the pulse oximetry screen.
 - Provide parents with a NBS blood spot and/or pulse oximetry brochure(s) & answer any questions they might have about the screen(s).



• Ensure the parents fill out a Refusal Form. Keep a copy for baby's record & fax a copy to the NBS Program using fax # 405-900-7556.

Refusal Form



Newborn Screening Program Religious Tenets and Practices Refusal Form

	Infant's Name:	Date of Birth:	Gender: N	M / F			
	Parent/Guardian's Name:		Medical Record #:				
	Street Address:		Apt/Unit #				
	City/State/Zip:		Phone #:				
	Place of Birth (check one):Hospital	_ Birthing Facility	Home Birth				
	Hospital/Facility Name:	Attending	Physician/Midwife:				
	Child's Dr/Planned Primary Care Provider:		Dr's Phone	#:			
	Type of Screen Refused:Newborn Blood (check any that apply & complete the corresponding se		Oximetry Screen	Hearing Screen			
	I, (Guardian's name), have been fully informed of the importance of newborn screening, and I understand that all newborns are required by law* to have the newborn screening tests performed. Although the benefits of newborn screening and the dangers of not being screened have been explained to me, I elect to refuse the newborn screening test(s) checked above for my child, (Infant's name), born on, on that such testing of my infant conflicts with my religious tenets and practices. My decision was made freely, and I accept the legal responsibility for the consequences of this decision. I have discussed the newborn screening tests with, my child's healthcare provider, and I understand the risks to my child if the newborn screen(s) are not completed.						
Refusal	I, (Guardian's name) screen test for are easily detected by testing a sm symptoms of these disorders sometimes do not a before symptoms become apparent. I have been i cause permanent damage to my child, including m	all blood sample from m ppear for several weeks nformed that these con	y baby's heel. I am awa or months, and irrever: ditions are treatable bu	sible damage can occur it if left untreated may			
Refusal	I, (Guardian's name), understand the congenital heart defects that the se oximetry test screen for can be detected by measuring the amount of oxygen in my baby's blood. I am aware that the is and symptoms of these defects sometimes do not appear for several weeks or months, and irreversible damage or the can occur if not identified early.						
Refusal	I, (Guardian's name) baby can hear sounds needed to listen and talk. It parents who can hear and there is no history of he the potential to interrupt speech, language, cogni	has been explained to rearing loss in their family	ne that most babies bo v. I understand that any	rn with hearing loss have			
	Print Parent/legal Guardian's Name Signa	ature of Parent/Legal Gu	ardian Date	/			

Transit Time

Prompt delivery of specimens to the Public Health Laboratory for testing can make all the difference.



Transit Time: What is it?



• "The time between the collection of a newborn screening specimen to its receipt at the OSDH Public Health Laboratory for testing."



Transit Time

• Guidelines:

- Specimens should be received at the OSDH Lab within **48 hours** from the time of collection.
- Oklahoma Law: OS 63 Sections 1-533 and 1-534

Delays in receiving the specimen


Delays in testing the specimen

Delays in diagnosis & treatment

Transit Time: Tips for Improvement

- Ensure everyone involved in NBS collection/handling knows about courier pick-up time, location, and importance.
- Do not batch specimens.
- Ensure the NBS is collected at 24 hr + 1 min of age & goes out with the courier as soon as possible after it has dried (\sim 3-4 hours of drying time).
- Set timelines and goals specific for your facility.
- Maintain a courier/transport log.
- Review transit time reports.
- Contact the PHL if the courier does not present to pick up the NBS specimens.

Oklahoma Birthing Hospitals by Courier Service





Public Health Laboratory Screening & Special Services
Oklahoma State Department of Health
Date created: 07/08/2023
Coordinate system: USA_Contiguous_Albers_Equal_Area_Conic
*44 birthing hospitals as of May 2023


Impact





Blood Spot Screening Statistics in Oklahoma

- In 2023 **50,294** dried bloodspot specimens were collected.
- 3 in 50 infants will have an abnormal screen that requires further testing.
- 1 in 450 infants will be identified with a disorder
- Specimens are kept by the OSDH lab for 42 days before being destroyed



^{**}LSD Disorders began screening in June 2021.



Newborn Screening WINS – True Story Timeline

Case Study from 2023

- Born on a Tuesday at 0121
- NBS collected Wednesday at 0150
- Specimen arrived at PHL Wednesday at 1900
- Preliminary critical result called to NBS follow up nurse on Thursday at 1210
- Baby was found to be still in hospital, in Mother/Baby unit. Mother/Baby nurse was notified, Genetic specialist notified and confirmatory labs were ordered. Feeding precautions initiated, decision made to delay discharge another night.
- NBS critical result finalized on Friday, called to NBS follow up nurse who promptly notified Mother/Baby staff and Genetics Specialist. Emergency management protocol in place, confirmatory labs are pending. Geneticist speaks with infant's parents at bedside for initial consultation. Treatment is initiated on Friday (3 days after birth), in anticipation of confirmatory testing results.
- Confirmatory lab results are finalized 8 days after birth. Ongoing care has been established with geneticist with plans for life long management in place.
- Delay of diagnosis or lack of treatment for this fatty acid disorder would have resulted in infant mortality.

Newborn Screening Loss – In Oklahoma in 2023

Case Timeline:

- Born on a Tuesday at 1904
- NBS collected Wednesday at 1951
- Specimen arrived at PHL Friday at 1900 (that Friday was a state recognized holiday)
- Friday 9 AM: NBS follow up team received word that infant was hospitalized in critical condition for R/O Sepsis, suspected seizures and was found to have elevated ammonia level. Geneticist requesting if NBS results were available? (Day 3 of life)
- Friday 4:12 PM: Infant passed on day 3 of life, screen and labs collected prior to passing indicated Citrullinemia to be cause of death

What do we learn from this family's experience?

- It was important to be able to trace the timeline, so that we knew that we had done all that we could to prevent this outcome for this family. (Courier logs are a part of this)
- Even when we control all the parameters within our scope, we will lose some infants to these disorders.
- We are compelled to address specimen quality and timeliness for every baby as if a disorder of this gravity hangs in the balance. We do have babies in Oklahoma diagnosed and living with Citrullinemia, due to early identification through newborn screening.

Pulse Oximetry Screening

CCHD – Critical Congenital Heart Disease



Critical Congenital Heart Disease (CCHD)

Screening began in 2014.

Screening is done by utilizing pulse-oximetry.

• Critical congenital heart defects are conditions that are present at birth and can affect the structure of a baby's heart and the way it works.

Pulse Oximetry Screening

Purpose:

• Screen **all** newborns between 24-48 hours of life with pulse oximetry to detect select defects related to critical congenital heart disease.

Rationale:

• Some newborns may appear healthy at first *despite* having a CCHD. Early detection and prompt treatment can prevent lifelong disability and early death.

Pulse-Oximetry Screening

- Simple and painless way to measure the amount of oxygen in the baby's blood.
- Congenital heart disease is the most common birth defect
- 1 in 110 infants will have a heart defect, 25% of those cases will have CCHD.
- Most affected will not have symptoms early on.
- Most will require surgery shortly after birth.

Normal Heart: Blood Flow




Image credit: CDC (2014)

- Blood from body tissues goes to the right side of the heart and enters the lungs, where the blood becomes oxygenated. The blood is then delivered to the left side of the heart, which is responsible for pumping the oxygenated blood out to the body in order to provide oxygenation to the body tissues. After being utilized, the deoxygenated blood is returned to the right side of the heart, and the cycle continues. Valves within the heart help to prevent backflow of blood during this process.
- Fetal openings between the atria, ventricles, and blood vessels begin to close shortly after birth.

Fetal-Neonatal Circulation

- The first breath of life leads to important changes in neonatal circulation:
 - Makes way for use of neonatal lungs (The lungs were not utilized in utero, as the placenta provided oxygenation to the fetus; after birth, however, an enormous amount of pressure is necessary in order for the newborn to close the diversions used to bypass the lungs in utero and instead allow for use of the lungs.)
 - Increased pressure change in the left side of heart compared to the right (The left side becomes the body's "pump") resulting in:
 - Closure of the Ductus Arteriosus (fetal opening between aorta and pulmonary artery)
 - Closure of the Foramen Ovale (fetal opening between the right and left atria)

❖ Failure of closure of fetal openings can result in complications

CCHD: Screening Targets & Symptomatology



CCHD Targets - Most likely detected by pulse oximetry screening

- Hypoplastic Left Heart Syndrome (HLHS)
- Pulmonary Atresia
- Tetralogy of Fallot
- Total Anomalous Pulmonary Venous Return
- Transposition of the Great Arteries
- Tricuspid Atresia
- Truncus Arteriosus
- * These heart defects lead to low levels of oxygen in the blood.

CCHD Targets - Potentially detected by pulse oximetry screening


- Double Outlet Right Ventricle (DORV)
- Ebstein's Anomaly
- Coarctation of the Aortic Arch
- Interruption of the Aortic Arch
- Single Ventricle
- Also potentially detected by pulse oximetry screening: other hypoxic cardiac or non-cardiac conditions.

CCHD: What to Watch For

Signs

- Cyanosis
- Tachypnea
- Increased work of breathing
- Swelling
- Tires easily during feeds
- Sweating
- Poor weight gain
- ❖ If at any time, the newborn should become symptomatic, the family should *immediately* take the baby to the closest emergency room for evaluation.

CCHDs in Oklahoma



Data provided by the Oklahoma Birth Defects Registry. Data does not reflect cases identified solely through pulse oximetry screening for CCHDs.

Pulse-Oximetry

The Screen and the Oximeter





Pulse Oximetry: Context

Who is screened?

- All newborns:
 - Must be calm & well; not crying
 - Warm extremities (temperature affects readings)
 - Skin clean & dry (dried blood affects readings)
 - Using room air; not on supplemental oxygen

When is screening performed?

- Healthy Newborn: Between 24-48 hours of life
- Sick Newborn: Between 24-48 hours of life
 - May delay if on supplemental oxygen
- **Before 24 hours:** higher risk for false positives (fetal-neonatal circulation transition not fully established)
- After 48 hours: delayed identification & treatment of affected newborns



The Pulse Oximeter

What is it?

- Screening tool: measures the percent of oxygen saturation of hemoglobin in the blood; and pulse rate
 - Simple
 - Painless
 - Non-invasive
 - Quick

The Pulse Oximeter

Oximeter Probe: 2 main parts

- light emitter
- Photodetector

Where is the probe placed?

- Right hand: pre-ductal measurement
- Either foot: post-ductal measurement

Points to Consider

- Pulse oximeter must be FDA approved (AAP, 2015)
- Regular calibration of the oximeter is required
- Pulse oximetry readings are averages
- Skin color and jaundice **do not** affect pulse oximetry readings
- **❖** Continuous pulse oximetry monitoring does not replace the pulse oximetry screen.

Screening How-to, Protocol and Guidelines





How is the Screen Performed?

- 1. Select site: right hand; either foot.
- 2. Place photodetector on outer aspect of hand/foot (under 4th-5th finger/toe).
- **3. Wrap** sensor tape around extremity.
- **4. Ensure** light emitter is **directly opposite** the photodetector.
- 5. If using a reusable sensor, secure the sensor using wrap recommended by vendor; **do not tape** or use hand to secure sensor to site.





Photo credit: Masimo 2011



Guidance for Screeners


Pulse Ox Dos

- If disposable, use a new, clean sensor; if reusable, clean between use
- Clean according to manufacturer recommendations
- Ensure newborn is calm and warm, not crying; encourage family involvement
- Ensure newborn skin is clean and dry
- Ensure no gaps between sensor and newborn's skin
- Light emitter and photodetector should be directly opposite of one another
- Use alongside physical examination
- Ensure pulse: no pulse, no oximetry!

Pulse Ox Don'ts

- Do not use an adult probe
- Do not tape pulse oximeter in place (use disposable wrap as indicated)
- Do not use your own hand to hold sensor in place
- Do not obtain reading from same extremity with blood pressure cuff
- Bilirubin lamps & surgical lights can affect accuracy of reading; cover pulse oximetry sensor with a blanket if such instruments are in use
- Do not use in isolation

Pulse Oximetry Screening Protocol



Screening Results

Negative Screen (Pass):

Oxygen saturation ≥ 95% in Right Hand and/or Left or Right Foot

AND

• Difference between the Right Hand and Left/Right Foot ≤ 3%

Positive Screen (Refer/Fail):

- Oxygen saturation < 90% in Right Hand or Left/Right Foot during any screen
- Oxygen saturation 90 94% for all 3 screens (1 hour between each screen)
- Difference between the Right Hand and Left/Right Foot > 3% for **all** 3 screens (1 hour between each screen)
- ❖If at any time, the newborn should become symptomatic, the family should immediately take the baby to the closest emergency room for evaluation.

Interpretation of Results

Negative = Pass

- Results are in-range
- Blood oxygen level WNL
- CCHD still possible (if symptomatic, a cardiac evaluation is warranted)
- Monitor baby's status:
 - Heart rate too fast/slow?
 - Energy overly sleepy/fussy/lethargic?
 - Appearance pale/blue skin?
 - Respiration too fast/slow?
 - Temperature cold to touch?
 - Feeding difficulties?

Positive = Fail/Refer

- Results are out-of-range
- Blood oxygen level is low
- High risk; not diagnostic
- Confirmatory procedures & referral for treatment are warranted

Interpretation of Results: CCHD Screening for Newborn without Cardiovascular

or Respiratory Distress

Oxygen Saturation

(O2 Sats; %)




Right Hand (RH)	Either Foot (F)			TT								
100	100	99	98	97	96	95	94	93	92	91	90	89% or lower
99	100	99	98	97	96	95	94	93	92	91	90	89% or lower
98	100	99	98	97	96	95	94	93	92	91	90	89% or lower
97	100	99	98	97	96	95	94	93	92	91	90	89% or lower
96	100	99	98	97	96	95	94	93	92	91	90	89% or lower
95	100	99	98	97	96	95	94	93	92	91	90	89% or lower
94	100	99	98	97	96	95	94	93	92	91	90	89% or lower
93	100	99	98	97	96	95	94	93	92	91	90	89% or lower
92	100	99	98	97	96	95	94	93	92	91	90	89% or lower
91	100	99	98	97	96	95	94	93	92	91	90	89% or lower
90	100	99	98	97	96	95	94	93	92	91	90	89% or lower
89% or lower	100	99	98	97	96	95	94	93	92	91	90	89% or lower

Pass	95% or higher in right hand (RH) or either foot (F) AND difference of 3% or less between RH and F.
Rescreen	90-94% in RH and F or difference of 4% or more between RH and F. Screen up to 3 times.
	89% or lower in RH or F (at any time)
Fail	OR
	3rd screen: 90-94% in RH and F <u>or</u> difference of 4% or more between RH and F.

Adapted from *Critical Congenital Heart Disease Screening with Pulse Oximetry in the Neonatal Intensive Care Unit* Lakshminsrusimha S., et al., e-Journal of Neonatology Research 2012(2)2: 96-101.



Reporting Results for CCHD: Filter Paper



Note: If parents refuse the pulse oximetry screen, provide them with a pulse oximetry brochure and answer any questions they might have about the screen. Ensure the parents fill out a Refusal Form; keep a copy for baby's record & fax a copy to the NBS Program using fax # 405-900-7556.

- Pulse Oximetry Screen: Check Only ONE
 - Pass
 - Fail
 - Not Performed
 - Refused
 - Echo



Reporting Results for CCHD - Pulse Oximetry Result Form

OKLAHOMA State Department of Health							
	e Department of Health						
Pulse Oximetry	Pulse Oximetry Screening Result Form						
Infant Information:							
Infant's Last Name:	Infant's First Name:						
Medical Record Number:	Attending Physician/Midwife:						
Date of Birth://	Birth Hospital:						
Mother's Last Name:	Mother's First Name:						
Pulse Oximetry Screening:							
Date of Screening://							
Age at Time of Screening:Days or	Age at Time of Screening:Days orHours						
Result:Pass/Negative	Result:Pass/NegativeFail/PositiveNot Performed						
Complete this section only if pulse oximetry so	Complete this section only if pulse oximetry screen was not performed:						
Reason pulse oximetry screen not performed:	Reason pulse oximetry screen not performed:						
Early Discharge	Early Discharge						
Screening Not Indicated due to							
Parent Refusal							
Screener's Name:	_						
Screener's Signature:							
paper. Original to infant's record, provide a copy Oklahoma State Department of Health, Newborn	Form to be utilized if pulse oximetry screening results were not documented on newborn screening filter paper. Original to infant's record, provide a copy to parent, and forward copy by fax or mail to: Oklahoma State Department of Health, Newborn Screening Program Coordinator, 1000 NE Tenth Street, Oklahoma City, OK 73117-1299, (405) 426-8220 or 1-800-766-2223; Fax (405) 900-7556.						
Health.Ok.gov							



Newborn Screening Contacts

Bloodspot, Pulse Oximetry, & Hearing Screening

Screening & Special Services 123 Robert S. Kerr Oklahoma City, OK 73102-6406 **Phone:** 1-405-426-8220

Toll Free: 1-800-766-2223

Fax: 1-405-900-7556

NewbornScreen@health.ok.gov

Public Health Laboratory

Newborn Screening Section Public Health Laboratory Service 4615 W. Lakeview RD Stillwater, OK 74075 Phone: 1-405-564-7750
Toll Free: 1-800-766-2223

Fax: 1-405-900-7611

Publichealthlab@health.ok.gov

Thank you for your time!

Questions?

