


Rising Tide of Congenital Syphilis

Anissa Lynch BSN, RN, Sexual Health Nurse
Consultant

Casey Von Woerkom MPH, STI Surveillance
Manager

OSDH Sexual Health & Harm Reduction
Service


OKLAHOMA
State Department of Health



Congenital Syphilis Oklahoma 2022



Health Equity in Practice



We recognize that systems, social and economic factors impact health.

Source: Institute for Clinical Systems Improvement, Going Beyond Clinical Walls: Solving Complex Problems (October 2014)




Syphilis Among Females, Oklahoma


| Diagnosis | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|------|------|------|------|------|
| Primary & Secondary Syphilis – Females | 144 | 244 | 338 | 417 | 481 |
| Primary & Secondary Syphilis – Pregnant Females | 11 | 19 | 25 | 41 | 42 |
| Total Syphilis - Females | 352 | 606 | 697 | 1135 | 1463 |
| Total Syphilis - Pregnant Females | 52 | 100 | 130 | 220 | 279 |
| Congenital Syphilis | 12 | 43 | 52 | 85 | 110 |



Congenital Syphilis - Rates of Reported Cases by Year of Birth and State and Territory, United States, 2022



Syphilis in Oklahoma: Congenital Cases and Rates Among Females, 2010-2022




Congenital Syphilis Cases in 2022, Maternal Information

- **110** CS cases in 2022
 - 43.6% (48/110) - Tulsa
 - 16.4% (18/110) - Oklahoma
- Maternal age ranged from 17-42 years
 - 31.8% - 30 to 34
 - 24.5% - 25 to 29
 - 24.5% - 20 to 24
- **63.6%** (70/110) had maternal drug use
- **74.5%** (82/110) had previous STI prior to syphilis diagnosis




Congenital Syphilis - Maternal Race and Disparities, 2022


Oklahoma Females, Aged 15-44 Years



CS Cases by Maternal Race/Ethnicity, 2022




Congenital Syphilis Cases in 2022, Timeline of Maternal Diagnosis




Congenital Syphilis Cases in 2022, by Prenatal Care and Screening

- **60.9%** (67/110) had prenatal care (PNC).
- Of those with PNC:
 - **76.1%** were tested for syphilis at their first appointment.
 - **55.2%** were tested at 28-32 weeks gestation
- **91.8%** (101/110) were tested for syphilis at delivery.



How Syphilis Spreads

- **Penetration:**
 - *T. pallidum* enters the body through direct contact with a syphilis sore during anal, vaginal, or oral sex.
 - From an infected mother to her unborn baby
- **Dissemination**
 - Travels via the circulatory system throughout the body
 - Can invade the nervous system (CNS) during any stage



Clinical Stages



- Syphilis goes through several stages
- Signs/symptoms and transmission risk differ by stage
- Stages may not progress linearly



Clinical Manifestations – Primary Stage

- First and most infectious symptomatic stage
- 1-5 weeks typical duration
- Serological tests may not be positive in early primary syphilis
- Painless sore ‘chancre’
 - Often singular
 - May not be seen
- Appears at site of infection (most commonly genital site)
- Will resolve without treatment but syphilis infection remains
- **It is possible for patients to have a non-reactive titer but still be infectious!**



Clinical Manifestations – Secondary Stage

- Occurs 0-6 weeks after primary chancre
- May overlap with primary stage
- 2-6 weeks typical duration
- Serological titer result usually highest during secondary stage
- Can be infectious
- May reoccur
- Symptoms resolve with or without treatment but syphilis infection remains

Symptoms Include:

- Rash (75 – 100%)
- Lymphadenopathy (50 – 86%)
- Malaise
- Mucous Patches (6-30%)
- Condyloma lata (10-2-%)
- Alopecia (5%)



Secondary Syphilis: Rash

- Rash shows through keratin layer
- May not be painful or itch
- Presents as different “types” of rashes – may resemble rashes caused by other conditions
- Palmar/Plantar and General Body Rash are hallmark characteristics of syphilitic rash



Clinical Stages of Syphilis

Secondary

Mucocutaneous Lesions
(Condyloma Lata)
Oral and genital
Mucous Patch



Clinical Stages of Syphilis

Secondary

Hair Loss
Patchy Alopecia



Early NP, NS; Unknown Duration/Late Syphilis

Early Non-Primary Non-Secondary

- Positive treponemal test result
AND
- Positive non-treponemal result
AND
- One of the following:
 - Negative test within last year
 - Signs/symptoms within last year
 - Sexual exposure to Primary, Secondary or Early NP NS with last 12 months
 - Sexual debut within last 12 months


Unknown Duration/Late

- Positive treponemal test result
AND
- Positive non-treponemal result
- Does not meet criteria for Primary, Secondary or Early NP NS



Clinical Manifestations – Neuro/ocular/otic Syphilis

- Syphilis can disseminate to the nervous system during any stage of syphilis.
- All clients should be assessed for neurosyphilis symptoms and tested for HIV.
- Referral for LP (VDRL screen) Should be done as soon as possible for clients with symptoms. Clients with reactive VDRL should be admitted for IV PCN Therapy.



Spirochetes in Neural Tissue



Uveitis


Symptoms and Manifestations include:

Visual Changes (Uveitis)
Hearing Loss
Headache
Poor Concentration/ Memory Loss
Personality or mood changes
Mental illness
Movement problems/loss of coordination
Paralysis
Meningitis
Muscle Contractions



Vertical Spirochete Transmission


Syphilis In Pregnancy



- ❑ Transplacental transmission of spirochetes from Maternal bloodstream to fetal bloodstream.
- ❑ Vertical Transmission rate is 90% if mother has untreated Primary or Secondary Syphilis. Fetus is at greatest risk during early stages of maternal infection.
- ❑ Treating maternal syphilis before the last month of pregnancy, decreases risk of congenital transmission by 98%.



Why do we care? Untreated syphilis is pregnancy will cause:



Early Congenital Syphilis

Clinical signs of congenital syphilis

| | |
|--------------------|-----|
| Osteochondritis | 55% |
| Snuffles | 40% |
| Rash | 40% |
| Anemia | 30% |
| Hepatosplenomegaly | 20% |
| Jaundice | 20% |
| Neurologic signs | 20% |
| Lymphadenopathy | 5% |
| Mucous patches | 5% |



Snuffles

From the delivering provider perspective, the placenta is often large, thick, and pale. The umbilical cord is edematous. Please send the placenta to pathology.



EARLY CONGENITAL SYPHILIS: CLINICAL MANIFESTATIONS: Newborn Rash



EARLY CONGENITAL SYPHILIS: CLINICAL MANIFESTATIONS

- Hepatosplenomegaly
- Anemia
- Thrombocytopenia
- Hydrops fetalis
- Pneumonia
- Nephrotic syndrome




Hepatosplenomegaly




**Flank Rash +
Hepatosplenomegaly**





Periostitis

Osteochondritis




Late Congenital Syphilis

- Hutchinson's Triad:
 - Interstitial keratitis
 - Hutchinson's teeth
 - Deafness (8th nerve)
- Short Maxilla
- High palatal arch
- Sternoclavicular thickening
- Frontal bossing
- Protruding mandible
- Clutton's joints
- Flaring scapulas
- Mulberry molars
- Saber shins



Fig. 1 : Dysmorphic facies with frontal bossing, prominent supraorbital ridges, absence of eyebrow and eyelashes, and a saddle nose



Who Needs Syphilis Screening

ALL PREGNANT FEMALES & Clients Who

- Have suspicious symptoms
- Recent partner(s) with a confirmed or suspected diagnosis
- History of or current Gonorrhea, Chlamydia, Syphilis or HIV diagnosis
- Multiple sex partners
- Homeless or hx of recent incarceration
- Hx of or current Drug Use
- Exchange sex for drugs/money/other needs
- **No Prenatal Care(Do not discharge until result is known!!)**



Syphilis Screening Recommendations in Pregnancy

CDC Recommendation is that all women who have delivered should have a current Syphilis test/result on file before being discharged!

**First Trimester
Or First PNV:
Everyone!!**



**Early 3rd trimester:
(28-32 wks.)**



**At Delivery:
EVERYONE!!**



**All Stillbirths
Miscarriage 20wks
or Greater.**



Management of Syphilis dx during second half of pregnancy should include sonographic evaluation for CS and additional HIV Screening!



Assessing, Diagnosing, & Treatment

- CDC Recommends a **Sexual history assessment** be done at each PNV for pregnant females diagnosed with STIs!
 - **Sexual practices** (oral, vaginal , rectal, condom use, substance use, sex work, dating apps etc.)
 - **Partners**
 - **New Partners, Multiple partners, anonymous partners**
 - **Treatment status of Partners!!!!**
 - **If partners have additional partners**
- History of syphilis
- Most recent serologic test for syphilis
- Known contact to an early case of syphilis
- Signs or symptoms of syphilis in the past 12 months
- **CURRENT sign/symptoms**
- **PREGNANCY Status**
- **SCREEN FOR HIV, GONORRHEA and CHLAMYDIA!!**
- **SCREEN for GC/CT at all reported sites of sexual contact (Vaginal, Oral, and Rectal)**



Physical Examination Questions

- Genitalia area
- Perianal area
- Oral cavity
- Skin of torso
- Palms and soles
- Neurologic examination

Neuro Syphilis Assessment

Change in or blurring of vision?

Recent eye pain or redness?

Spots or distortion in vision?

Double vision?

Light hurting eyes?

New weakness in arms, legs, or face?

New headache unlike usual headaches?

Stiff neck?

New/recent hearing loss?

New/recent ringing of ears?

Ocular injection

Photophobia

Nuchal rigidity

Facial palsy



Laboratory Testing

Diagnosis of syphilis requires a positive test from **both** of these categories with the exception of early primary syphilis:

Non – Treponemal


- **RPR**
- VDRL
- Detects non-specific antibodies caused by inflammation
- Quantitative if reactive - measured in titers (*helps determine past/present infection or reinfection and treatment efficacy*)
- **May be reactive for life post treatment (Low Serofast)**
- Use this to determine if patient is cured or has treatment failure or re-infection

Treponemal

- **TP-N, TPPA, MBIA**
- FTA-ABS, EIA/CIA
- Detects specific antibodies against *T. pallidum*
- **Likely remains positive forever in most(85%)**
- Qualitative (Reactive/Non-Reactive)
- Can be screening or confirmatory



Serologic Titers of RPR and VDRL



Persons who have signs or symptoms that persist or recur and those with at least a fourfold increase in non treponemal test titer persisting for >2 weeks likely were re-infected or experienced treatment failure. (CDC Treatment Guidelines,2021).



Who Needs Treatment



All symptomatic clients
All asymptomatic clients with a reactive titer whose Titer/treatment history can't be confirmed.



All pregnant females who are:

- **Symptomatic,**
- **Reactive serology with no symptoms or and No history of syphilis**
- **Has partner(s) with a confirmed diagnosis.**



All partners to a confirmed case of syphilis-

Confirmed case= Partner has primary or secondary symptoms, partners reactive titer status is k
DIS have confirmed partner is infected and untreated.




Image Credit: Louisiana Department of Health



Treatment of Syphilis in Pregnancy

- The only treatment for syphilis in pregnancy is penicillin. There are no available alternatives.
- Pregnant women should be treated with the penicillin regimen appropriate for their stage of infection.
- Pregnant women with penicillin allergy require desensitization in a hospital setting due to the risk for serious IgE – mediated hypersensitivity reactions (CDC, 2021).
- Treating Partners of pregnant females is crucial to preventing reinfection before delivery!



Syphilis Treatment During Pregnancy

Draw Blood for Syphilis titer on first day of treatment

P & S symptoms and Early Latent=
Treatment x's 1 dose




Total dose Bicillin 2.4mu

Pregnant P & S symptoms and Early Latent=
Treatment x's 2 doses at 1 week intervals

Day 1



Day 7



Total dose Bicillin 4.8mu

Late Latent and unknown duration (NO symptoms)=
Treatment x's 3 doses at 1 week intervals


Day 1



Day 7



Day 14



Total dose Bicillin 7.2mu

Bicillin is the only approved regimen for Pregnant Females. Patients reporting PCN allergy must be desensitized and treated with BIC!!!



Bicillin Treatment in Pregnancy: Time Between Doses for Latent Syphilis

- Adherence to 7 day interval between doses in pregnancy is necessary
- 40% of pregnant woman are below treponemicidal levels after 9 days
- Pregnant females must Restart entire series (3 weekly doses) if dose is missed (interval >7 days)
- **Treatment must be initiated and completed > 30 days before delivery to be considered adequate.**
- Encourage Abstinence or Condom use until patient and partner(s) are treated.



Jarisch-Herxheimer Reaction



Acute febrile illness following penicillin injection

Usually occurs within a few hours after Bicillin injection



Dying treponemes release endotoxins faster than body can clear them



Can occur at any stage

Most common during early syphilis when treponemal load is higher



Can cause exacerbation of syphilis symptoms and uterine contractions



Follow Up Serology Titers Post Treatment

- Primary/Secondary/ Early Latent Syphilis- 6 & 12 months*
- Latent/Unknown Duration - 6, 12,& 24 months*
- Clients Living with HIV- 3, 6, 9, 12, & 24 months
- Pregnant females diagnosed and treated at or before 24weeks- 8 weeks post treatment
- Pregnant females diagnosed and treated after 24 weeks- Follow titers should be repeated at delivery**

* Clients with risk factors, practices, or behaviors pose an increased risk for syphilis acquisition can be screened more frequently.

****Titers should be repeated sooner in pregnant females if reinfection or treatment failure is suspected or client is living with HIV****



Barriers To Treatment

- Insurance
- Transportation
- Patient/Provider follow through
- **Bicillin Shortage! (Beginning in January 2023, the U.S. is having a national Bicillin shortage. Bicillin is now prioritized for Pregnant Females and Clients living with HIV. All non-pregnant individuals are being treated with Doxycycline regimens).**



Nurses Responsibility in Syphilis Staging and Treatment



- Take a Thorough Sexual History! Including assessment of risk behaviors and treatment of sex partners to assess for reinfection (CDC STI Treatment Guidelines 2021)
- **Ensure Physical Exam and Specimen Collections are done.**
- Accurate Staging of Syphilis symptoms. (Always review chart for titer hx.)
- **ASK QUESTIONS** about PCN allergy!
- Provide Client Centered Prevention and Follow-UP Education
- Ensure new syphilis case is put into PHIDDO **within 24 hours** of receipt of result!!
- Collaborate with DIS on referrals(desensitization, LP for neurosyphilis exam), information on titer history and staging clients.
 - Call or **email** your DIS when you have symptomatic client in clinic(Include all pertinent details(stage/treatment pregnancy status and good contact info for client)
 - Prioritize getting pregnant females and partners into clinic for exam and treatment!!
- Contact SHHR Nurse Manager or Nurse Consultants if you have questions about treatment or circumstance that is not in protocol!




Disease Intervention Specialist (DIS)

DIS interview patients and contacts to gather information concerning infections or exposure to HIV and Syphilis. Consults with private physicians to stimulate case reporting, obtain information on treated cases, provide information on diagnostic and treatment techniques, and link clients and exposed partners to facilities for treatment.

Collaboration for the exchange of clinical information on clients between providers and DIS is critical to ensure clients are screened and treated appropriately.

Disease Intervention Specialist (DIS) Workforce



DIS Responsibilities



- Interview clients with a newly reactive syphilis titer, symptom, contact or re-infected.
- Find and interview partners who have been a contact to syphilis (all pregnant females, and partners of OC with a titer $\geq 1:16$).
- Draw lab specimens.
- Assist with Transporting clients to CHD if needed.
- Stage syphilis based on the information obtained during client interview.
- Assist with tracking titers and treatment history of patient.
- Collaborate with CHD and nursing staff to refer client to care, assist with staging and symptom identification.
- Provide Client Centered Prevention information.



Things to remember for Management of Syphilis in Pregnancy

- Initial Staging can change based on provider examination or DIS investigation.
- If client has a history of syphilis, The RPR will determine if client is re-infected
- TP-N & TP-PA, FTA, MBIA will always be positive after initial infection
- Ask specific questions about reported PCN allergy to determine if client can still be treated with BIC.(When? What type of reaction? Was medical intervention needed?)
- **Test all clients for HIV, Gonorrhea and Chlamydia!**



Things to remember for Management of Syphilis in Pregnancy

- Always treat symptomatic clients regardless of if test result is non-reactive or not back yet!!! Remember that titers are sometimes delayed to rise in primary syphilis.
- DO NOT delay treatment or referral if your DIS has not called you back! Initiate treatment, create PHIDDO case and email your DIS the information!
- Treat for appropriate stage of syphilis. Missed Doses greater than 14 days(non-pregnant clients) and 9 days for pregnant females or require client to start treatment regimen over.
- **ALWAYS ASSESS FOR NEURO/OCULAR/OTIC SYPHILIS.** Give treatment with BIC according to Stage and refer to ER for immediately for CSF-VDRL Screen. **BIC WILL NOT TREAT NEURO SYPHILIS!!**
- **DO NOT GIVE BIC IN A DELTOID MUSCLE!!!!** This is not best practice, extremely painful for client. BIC is poorly absorbed due to small muscle mass, and can cause severe muscle damage including necrosis!!!!



How Oklahoma Providers Can Help

- Become Familiar with Syphilis Symptoms and CDC Treatment recommendations
- Communicate titer results and treatment status between OB and PEDS provider.
- Work with OSDH DIS, Maternal FSS and Nurses and County Health Departments to ensure patient and partner(s) are staged and treated appropriately.
- O.S. § 1-502.2 protects health care providers to share medical or epidemiological information to health professionals , state agencies or district court within the continuum of care for diagnosing and treating communicable diseases without a written ROI.



How Oklahoma Providers Can Help

- Incorporate comprehensive sexual history questions into EMR platform.
- Consider creating internal CS nurse/physician case management committee to keep track and contact with pregnant clients diagnosed with syphilis to ensure they follow through with treatment, partner treatment and to work with in house lab to improve time of getting syphilis results.
- Incorporate in house guidelines for PCN desensitization, oral challenge, or skin testing to ensure Pregnant females with PCN allergy can be treated with Bicillin as soon as possible.
- Complete Reporting of Necessary Variables (Report Syphilis to OSDH within 24hours!)
 - Demographics, Pregnancy Status
 - Signs/Symptoms and Treatment
 - Details what would help guide intervention



How Oklahoma Providers Can Help

- Consider joining Congenital Syphilis Review Board and Task Force. Contact Dawn.Kluesner@health.ok.gov
- Consider enrolling RNs and APRNs into STI Academy. Contact Anissa.Lynch@health.ok.gov
- Consider keeping condoms in triage/ED/offices to provide to symptomatic clients or clients with diagnosis. Contact Sexual Health and Harm Reduction Service for information on how to obtain free condoms.
- Build Rapport with DIS and Local County Health Departments! CHDs will treat pregnant females for free.



SHHR Contact Info

Phone:
405.426.8924

Fax:
405.900.7585

Webpage:
SHHR.health.ok.gov

Email:
Condoms@health.ok.gov
HIVSTDTESTS@health.ok.gov

Rapid Start/PrEP
RapidStart@health.ok.gov

Need condoms for events? Email the condoms@health.ok.gov email address. Please include Name/date of event, Name and shipping address, and number of condoms/lube requested. Please submit request at least 2-3 weeks before event.



For Additional STI Information

CDC Sexually Transmitted Infections Treatment Guidelines 2021
<https://www.cdc.gov/std/treatment-guidelines/default.htm>

National Network of STD Clinical Prevention Training Centers
www.STDCCN.org

Sexual Health and Harm Reduction Service (405)-426-8400 <https://shhr.health.ok.gov>

App available on Apple
and Android Devices



iOS and android Users
Launch Chrome or Safari on your
phone or tablet
Navigate to STI mobile app (cdc.gov)
Tap the Menu icon or Share Icon
Tap Add to Home Screen.



CDCS
TD



@CDCSTD



Amber Rose, MS
APM Surveillance and Analysis
Sexual Health and Harm Reduction Service
amberer@health.ok.gov
405.973.8461

Anissa Lynch BSN, RN
STI Nurse Consultant | Rapid Start Program
Manager
Anissa.Lynch@health.ok.gov **405.540.2708**

Casey Van Woerkom MPH,
STI Surveillance Manager
CaseyV@health.ok.gov **405.426.8722**