



Preeclampsia & Eclampsia: Hypertensive Disorders of Pregnancy

Evelyn Radichel, MS, RN
Melissa Warde, MHA, CAPM



Goals and Objectives

01

Define clinical criteria for:

- Gestational hypertension
- Preeclampsia
- Eclampsia
- HELLP syndrome

02

Describe alterations of maternal physiology in preeclampsia

03

Prioritize care of the eclamptic patient during a convulsion



***How many types of
hypertension can a woman
have?***


Hypertensive Disorders of Pregnancy: 4 Classifications

- 1) Chronic Hypertension
- 2) Chronic Hypertension + Preeclampsia
- 3) Gestational Hypertension
- 4) Preeclampsia
 - a) Severe Preeclampsia
 - a) Eclampsia
 - b) HELLP syndrome


2022 OSDH: 8.8% or 1 in 11 pregnant Oklahomans experienced a hypertensive disorder of pregnancy.



Chronic Hypertension vs. Gestational Hypertension



Spectrum of Hypertensive Disorders of Pregnancy



Definitions: 140/90 = Hypertension

160/110 = Severe Hypertension

- Chronic (preexisting) Hypertension
 - Onset prior to 20th week gestation
 - May continue through 12 wks PP
 - BP \geq 140 systolic or \geq 90 diastolic
 - BP \geq 160 systolic or \geq 110 diastolic = **Severe**
- Gestational Hypertension
 - Occurring after 20 weeks gestation in a previously normotensive woman
 - Resolves by 12 wks PP
 - BP \geq 140 systolic or \geq 90 diastolic
 - BP \geq 160 systolic or \geq 110 diastolic = **Severe**


Preeclampsia

*Syndrome defined by hypertension & proteinuria
(proteinuria is not a requirement for diagnosis)*

Gestational Hypertension
(140/90)

+

Any of these symptoms



- Proteinuria
 - $\geq 300\text{mg}/24\text{ hr.}$ or
 - Protein/Creatinine ratio ≥ 0.3
 - Dipstick $\geq 2+$
- Cerebral or visual symptoms
- Epigastric or right upper quadrant pain
- Oliguria $< 500\text{ ml.}$ over 24 hours
- Thrombocytopenia $< 100\text{K plt.}$
- Elevated liver enzymes (twice normal)
- Pulmonary edema
- New onset headache - unresponsive to medications

Severe Preeclampsia

- Systolic BP \geq 160, or diastolic BP \geq 110
 - *Elevated SBP is better indicator of stroke than DBP
 - (95.8%) women with systolic BP > 160mm Hg →stroke
 - (12.5%) women with diastolic BP > 110mm Hg →stroke
- Severe headache and vision changes
- Persistent epigastric or right upper quadrant pain
- Oliguria < 500 ml. over 24 hours
- Nausea & vomiting
- Thrombocytopenia
- Elevated liver enzymes (twice normal)
- Development of Eclampsia
- Development of HELLP syndrome
- Pulmonary edema





Risk Factors for Preeclampsia

- Nulliparity
- Multifetal gestations
- Preeclampsia in a previous pregnancy
- Chronic hypertension
- Pregestational diabetes
- Gestational diabetes
- Thrombophilia
- Systemic lupus erythematosus
- Prepregnancy body mass index greater than 30
- Maternal age 35 years or older
- Antiphospholipid antibody syndrome
- Kidney disease
- Assisted reproductive technology
- Obstructive sleep apnea



Maternal Complications

- Stroke from cerebral hemorrhage
- Placental abruption
- Eclamptic seizures
- Cerebral Edema
- Liver hematoma/rupture
- Pulmonary edema
- Acute renal failure
- Hemorrhage/DIC
- Cardiomyopathy



Fetal Complications

- IUGR
- Premature birth
- Fetal intolerance to labor
- Hypoxia
- Death



***What is the process behind
the disease?***



Pathophysiology

Failure of normal physiologic adaptations to pregnancy


Normal Pregnancy

- ↑ plasma volume
- ↓ vascular resistance
- ↑ renal blood flow

Preeclampsia

- ↓ plasma volume
- ↑ vascular resistance
- ↓ renal blood flow

Probably multiple etiologies cause the syndrome
– “Disease of Theories”



Non-pregnant

Normal pregnancy

Pre-eclampsia/IUGR


Spiral arteries
Normal pregnancy
Dilates →
increased bloodflow


Spiral arteries
Preeclampsia:
Fibrous → narrow
Less bloodflow

Pathophysiology

- Stage 1: Poor placentation
 - Incomplete invasion of spiral arterioles results in
↓ uteroplacental blood flow
- Stage 2: Inflammation
 - The ischemic placenta induces widespread endothelial cell damage and maternal systemic inflammatory response

Alteration in Systems Function





***What are best practices to
care for these women?***



5 Management Objectives

1. Recognize the situation (signs & symptoms)

- ① 2 elevated BP within 15 min. → notify physician
- ② Initiate anti-hypertensive treatment ASAP

2. Control BP with antihypertensive agents

↓ Arterial spasm to prevent vascular injury to brain, kidneys, and heart

Diastolic not below 90: placenta needs adequate perfusion

3. Prevent or control seizure activity

Magnesium Sulfate infusion

4. Delivery of fetus

Consider GA and delivery route

5. Postpartum surveillance

3-10 day follow-up in provider office (72 hours)

Delivery Timing Considerations

37 weeks – deliver

34 weeks – deliver after maternal stabilization AND

- Antenatal steroids –Betamethasone
- Deliver in 48 hours

Deliver as soon as maternal stabilization with following complications:

- Fetal reasons: concerning FHR pattern, poor Doppler studies...
- Abruption
- Pulmonary edema
- Eclampsia –stabilized
- DIC
- Persistent/worsening symptoms

Medications for Severe Hypertension

- Treatment of critically elevated BP with either
 - IV labetalol
 - IV hydralazine
 - Oral nifedipine

⊘ Oral labetalol onset is slow and peaks at 1-4 hours. Expected to be less effective.


Initial Management Begins With	Initial Dose	Next Dose ^a	Next Dose ^a
IV Labetalol (Beta blocker) Side effects: low heart rate bronchoconstriction	<ul style="list-style-type: none"> • Labetalol 20 mg IV for more than 2 min • Check BP in 10 min 	<ul style="list-style-type: none"> • Labetalol 40 mg IV for more than 2 min • Check BP in 10 min 	<ul style="list-style-type: none"> • Labetalol 80 mg IV for more than 2 min • Check BP in 10 min
IV Hydralazine (Arteriolar) Side effects: tachycardia, hypotension, flushing, headache	<ul style="list-style-type: none"> • Hydralazine 5 mg or 10 mg IV for more than 2 min • Check BP in 20 min 	<ul style="list-style-type: none"> • Hydralazine 10 mg IV for more than 2 min • Check BP in 20 min 	<ul style="list-style-type: none"> • Labetalol 20 mg IV for more than 2 min • Check BP in 10 min
Oral nifedipine (Ca Chanel blocker) Side effects: low BP low HR+ dizziness	<ul style="list-style-type: none"> • Immediate release Nifedipine capsules (10 mg orally) • Check BP in 20 min 	<ul style="list-style-type: none"> • Immediate release Nifedipine capsules (20 mg orally) • Check BP in 20 min 	<ul style="list-style-type: none"> • Immediate release Nifedipine capsules (20 mg orally) • Check BP in 20 min

Contraindicated
In asthma

If poor results after 3 doses
move on to another
antihypertensive
Doctor should obtain consult

Magnesium Sulfate Therapy

- Use extreme caution when administering drug
 - High alert drug (ISMP)
- Monitor output- renal excretion
- Monitor deep tendon reflexes (DTRs)
- Monitor respirations
- Monitor LOC



Magnesium Toxicity

- Therapeutic: 4-8mg/dl
- Loss of DTR's: 9-12mg/dl
- Respiratory arrest/muscle paralysis: 12-18mg/dl
- Cardiac arrest: 25-30mg/dl

Renal excretion – beware of DM and other ↓ renal function.



Magnesium Toxicity



Antidote

Calcium Gluconate 10%

- 1g/10ml IV over 3 minutes



Airway and ventilatory support as needed



O2 and suction set up and ready

Management of Preeclampsia

- Magnesium Sulfate is drug of choice
 - Acts at neuromuscular junction to produce muscular relaxation.
 - Small vessel vasodilation.
- Loading dose of 4-6g over 15-30 minutes
- Followed by maintenance dose of 2-3 g per hour

Magnesium Sulfate is NOT an antihypertensive medication.

Magnesium Sulfate Therapy

- Increases Ca^+ excretion
- Crosses placenta – be prepared for a lethargic infant – may require resuscitation
- Increases possibility of postpartum hemorrhage
 - **NO Methergine**, **Cytotec** preferable.

Eclampsia

New onset of convulsions and/or coma in a woman with signs of preeclampsia

50%
antepartum

25%
intrapartum

25%
postpartum



Mechanism: cerebral edema, ischemia, hemorrhage or vasospasm

Complications of Eclampsia

Placental
abruption

Pulmonary
edema

Aspiration
pneumonia

Cerebral
hemorrhage

Renal tubular
necrosis



Liver rupture

Retinal
detachment

Dissemination
intravascular
coagulation (DIC)

Cerebral Hemorrhage

****Cerebral hemorrhage is a common autopsy finding**



Section of brain at autopsy

Eclampsia

Indicators of Seizure

- Headache – most common indicator
- Visual disturbances – i.e. seeing spots, loss of vision, etc
- Epigastric Pain – RUQ pain
- **NO INDICATORS** – just suddenly occurs

Management of Eclamptic Convulsion

Life-threatening emergency requiring immediate action.

- Prevent injury to woman
- Maintain airway
- Magnesium Sulfate to control convulsion



Management of Eclamptic Convulsion

Magnesium Sulfate Regimen



4-6g loading dose given over 15 min followed by 2-3g/hr maintenance – onset of action is immediate.



If seizure reoccurs, may administer another 2g over 5 min.



If seizure continues or reoccurs may sedate, intubate and ventilate.

Hematoma following seizure



Post Seizure Care

Do not leave patient alone

- Maintain quiet environment
- O₂ at 8-10L/min. per mask
- VS Frequently
- Monitor SaO₂ : $\geq 95\%$



Post Seizure Care

- Maintain magnesium infusion as ordered
- Labs and chest x-ray
- If undelivered:
 - Monitor FHR and uterine activity
 - Consider route of delivery
- Designate someone to keep family informed

Post Seizure Care


- Observe for signs of:
 - Abruption if undelivered
 - Pulmonary edema
 - Neurologic deterioration
 - Oliguria - output < 30cc/hr
 - Signs of magnesium toxicity:
 - ↓ DTR's, respirations < 12/min.

Eclampsia – Documentation of Occurrence



- Time seizure began
- Duration of seizure
- Care provided
- Maternal and fetal responses
- Duration of postictal phase
- Duration of unconsciousness (if unconscious)

HELLP Syndrome



- Hemolysis

- Abnormal peripheral blood smear- schistocytes & burr cells
- ↑ bilirubin

- Elevated Liver enzymes -2X upper limits of normal


- LDH > 600 IU/L
- ALT > 70 IU/L

- Low Platelets

- Thrombocytopenia < 100,000mm³
- Severe < 50,000mm³

HELLP Syndrome

- Frequently does not present with classic preeclamptic symptoms of hypertension & proteinuria.
 - Malaise - 90%
 - R ↑ quad. Pain - 65%
 - N/V - 50%
 - Worsening edema
 - Abdominal, flank or shoulder pain
 - Hematuria
 - Hypoglycemia



HELLP Syndrome

May be misdiagnosed as:

- Gall bladder disease
- Gastritis
- Appendicitis
- Pyelonephritis
- Acute Fatty Liver of Pregnancy

Any pregnant woman presenting with these symptoms should have:

- CBC with peripheral smear
- Liver Function Tests

Maternal Morbidity and Mortality

- Cardiovascular disease is the most prevalent preventable cause of maternal mortality in developed countries
- In the US, one-third of pregnancy-related deaths are attributed to cardiovascular conditions, and approximately 60% of these deaths are deemed preventable
- Maternal stroke is an infrequent but debilitating complication of pregnancy and it accounts for at least 7.7% of pregnancy-related deaths in the US
- Specific pregnancy-related conditions often trigger maternal stroke, many of which are potentially preventable
- The risk of stroke among pregnant and post-partum women is ~3 times increased compared with non-pregnant women of similar age

Maternal Stroke

Table 1:

Rates of maternal stroke across different nations ←

Study	Country	Enrollment period	Pregnant/post-partum women, n	Incidence of maternal stroke (per 100,000)	Mean age of women with stroke, years
Elgendy et al. ¹⁶	USA	2007–2015	37,360,772	45	30
Liu et al. ¹⁷	Canada	2003–2016	3,907,262	13.4	NR
Yoshida et al. ¹⁹	Japan	2012–2013	2,115,949	10.2	32.2
Sharshar et al. ⁸	France	1989–1992	669,680	4.6	30.6
Bashiri et al. ²⁰	Israel	1988–2004	173,803	9.2	35.5
Liang et al. ²¹	Taiwan	1992–2004	66,781	47.9	30.1
Prabhu et al. ¹²	India	2006–2008	39,211	66	22





Mechanisms of Pregnancy-Associated Stroke




Figure 3:

Potential mechanisms of maternal stroke.

Summary of possible mechanisms of maternal stroke.


PFO= patent foramen ovale; RCVS= reversible cerebral vasoconstriction syndrome



***What do the facts tell us
about Severe Maternal
Morbidity (SMM)?***

Healthcare Provider Contributing Factors

Highest rates of preventability:
Hemorrhage 70%
Preeclampsia 60%



Health care provider factors were the most common type of contributor identified for all five leading causes of death and were particularly common for preeclampsia and hemorrhage, consistent with their higher degree of preventability. The most common health care provider factor was delayed response to clinical warning signs followed by ineffective care.


Main et al. 2015. Pregnancy-related Mortality in California
Obstetrics & Gynecology



Poor Management Outcomes



- *Critical symptoms not recognized (Denial)*
- *Delayed diagnosis*
- *Delayed treatment*
- *Inadequate treatment*
- *Assumption that delivery cured preeclampsia*
- *Discharge without timely follow-up*



“The best way to diagnose preeclampsia is to listen to your patients.” ~ Dr. Baha Sibai

Most Common Preventable Errors

See It!

- Failure to adequately control blood pressure in hypertensive women

Believe It!

- Failure to adequately diagnose and treat pulmonary edema in women with preeclampsia

Treat It!

- Failure to pay attention to vital signs following birth
- Hemorrhage following cesarean birth

Preeclampsia related



OKLAHOMA PERINATAL
NURSE FORUM

Knowledge is Power:

OB Emergencies in the ED

Hear Her Campaign:

<https://youtu.be/tXBfzmgJFCU>



Solutions to Reduce the Impact of Maternal Hypertension

Table 4:

Call for action- Clinical practice and healthcare settings

1. Improved prenatal patient education of cardiovascular risk factors, symptoms of cardiovascular complications of pregnancy, and the importance of long-term preventative care among reproductive age women.
2. Implementation of multidisciplinary healthcare team education and maternal stroke toolkits to improve recognition of cardiovascular complications and standardization of maternal healthcare delivery.
3. Reduction of socioeconomic disparities in maternal cardiovascular outcomes, through increased access to healthcare coverage for pregnant and postpartum women, increased access to maternal healthcare in rural areas, and efforts to address systemic racism.
4. Targeted efforts to reduce knowledge gaps in maternal cardiovascular health, through increased funding for maternal cardiovascular research and increased inclusion of pregnant and postpartum women in clinical trials.

Discharge Planning/ Teaching

- Any patient treated for hypertension or preeclampsia **f/u in 3-7 days**
- *Delivery is not a cure-* Preeclampsia can occur up to 6 weeks PP
- Teach symptoms of Preeclampsia to **all patients**

Why should you care?

What should you pay attention to?

What you should you do if you have any signs

Preeclampsia Foundation

Ask Your Doctor or Midwife

Preeclampsia

What Is It?

Preeclampsia is a serious disease related to high blood pressure. It can happen to any pregnant woman.

Risks to You

- Seizures
- Stroke
- Organ damage
- Death

Risks to Your Baby

- Premature birth
- Death

Signs of Preeclampsia



Stomach pain



Headaches



Feeling nauseous, throwing up



Seeing spots



Swelling in your hands and face



Gaining more than 5 pounds in a week

What Should You Do?

Call your doctor right away. Finding preeclampsia early is important for you and your baby.

For more information go to www.preeclampsia.org

Copyright © 2010 Preeclampsia Foundation. All Rights Reserved.

AWHONN Post-Birth Warning Signs Handouts

POST-BIRTH
Warning Signs
Education Program
- AWHONN

SAVE YOUR LIFE:

Get Care for These POST-BIRTH Warning Signs

Most women and postpartum people who give birth recover without problems. **But anyone can have a complication for up to one year after birth.** Learning to recognize these POST-BIRTH warning signs and knowing what to do can save your life.

Trust your instincts.
ALWAYS get medical care if you are not feeling well or have questions or concerns.

<p>Call 911 if you have:</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Pain in chest <input type="checkbox"/> Obststructed breathing or shortness of breath <input type="checkbox"/> Seizures <input type="checkbox"/> Thoughts of hurting yourself or someone else
<p>Call your healthcare provider if you have: (you only need one sign) <small>(If you can't reach your healthcare provider, call 911 or go to an emergency room)</small></p>	<ul style="list-style-type: none"> <input type="checkbox"/> Bleeding, soaking through one pad/hour, or blood clots, the size of an egg or bigger <input type="checkbox"/> Incision that is not healing <input type="checkbox"/> Red or swollen leg, that is painful or warm to touch <input type="checkbox"/> Temperature of 100.4°F or higher or 96.8°F or lower <input type="checkbox"/> Headache that does not get better, even after taking medicine, or bad headache with vision changes

Tell 911 or your healthcare provider:

"I gave birth on _____ and
(Date)
I am having _____"
(Specific warning signs)



Scan here to download this handout in multiple languages.

These post-birth warning signs can become life-threatening if you don't receive medical care right away because:

- **Pain in chest, obstructed breathing or shortness of breath** (trouble catching your breath) may mean you have a blood clot in your lung or a heart problem
- **Seizures** may mean you have a condition called eclampsia
- **Thoughts or feelings of wanting to hurt yourself or someone else** may mean you have postpartum depression
- **Bleeding (heavy)**, soaking more than one pad in an hour or passing an egg-sized clot or bigger may mean you have an obstetric hemorrhage
- **Incision that is not healing, increased redness or any pus** from episiotomy, vaginal tear, or C-section site may mean an infection
- **Redness, swelling, warmth, or pain in the calf area** of your leg may mean you have a blood clot
- **Temperature of 100.4°F or higher or 96.8°F or lower**, bad smelling vaginal blood or discharge may mean you have an infection.
- **Headache (very painful), vision changes, or pain in the upper right area of your belly** may mean you have high blood pressure or post birth preeclampsia



This program is supported by funding from Merck through Merck for Mothers. Merck for Mothers is known as MSD for Mothers outside the United States and Canada.

AWHONN thanks Kenvue for commercial support of the translations of this handout.

©2023 Association of Women's Health, Obstetric, and Neonatal Nurses. All rights reserved. Unlimited print copies permitted for patient education only. For all other requests to reproduce, please contact permissions@awhonn.org.

16005

**OKLAHOMA PERINATAL
NURSES FORUM**

Prognosis and Long Term Effects of Eclampsia

- Women with severe preeclampsia ↑ risk of developing cardiovascular disease later in life
 - Hypertension, Ischemic heart disease, stroke, dementia
 - *Many women are not aware of the long-term complications associated with preeclampsia*
- Preeclampsia with preterm delivery is a strong risk factor for CV disease (AHA)
- Conclusion of all is that pregnancy may be a screening test for chronic hypertension and CV disease

Long Term Maternal Outcomes

- Chronic hypertension (4 fold)
- Ischemic heart disease (2 fold)
- Stroke (2 fold)
- Venous thromboembolism (2 fold)
- All-cause mortality (1.5 fold)
- Recurrent preeclampsia

Prevention of Preeclampsia

ACOG recommends the use of low-dose aspirin (81 mg/day), initiated between 12 and 28 weeks of gestation, for the prevention of preeclampsia in individuals with 1 high-risk factor, or 2 or more moderate-risk factors.

TABLE 2

Risk factors and recommendations for preeclampsia⁶

Risk level ^a	Risk factors	Recommendation
High ^b	<ul style="list-style-type: none"> History of preeclampsia, especially when accompanied by an adverse outcome Multifetal gestation Chronic hypertension Pregestational type 1 or 2 diabetes Kidney disease Autoimmune disease (ie, systemic lupus erythematosus, antiphospholipid syndrome) Combinations of multiple moderate risk factors 	Recommend low-dose aspirin if the patient has ≥ 1 of these risk factors.
Moderate ^c	<ul style="list-style-type: none"> Nulliparity Obesity (ie, body mass index > 30) Family history of preeclampsia (ie, mother or sister) Black individuals (due to social rather than biological factors)^d Lower income^d Age ≥ 35 y Personal pregnancy factors (eg, infant of low birth weight or small for gestational age, previous adverse pregnancy outcome, > 10-year pregnancy interval) In vitro conception 	<ul style="list-style-type: none"> Recommend low-dose aspirin if the patient has ≥ 2 moderate risk factors. Consider low-dose aspirin if the patient has 1 of these risk factors.^d
Low	Previous uncomplicated full-term delivery	Do not recommend low-dose aspirin.



^a Includes only risk factors that can be obtained from the patient's medical history.

***Case Review:
What would you do?***

24 yo G1P0 at 34 weeks

- Presented to hospital at 2300
- Reports decreased fetal movement and headache
- BP 165/105 Right Arm
- No proteinuria
- Patient to left side
- Reactive NST
- BP now 150/100
- Given Vicodin for HA... HA improved
- DC home

24 yo G1P0 at 34 weeks

- Presented to hospital at 2300
- Reports decreased fetal movement and headache
- BP 165/105 BP not treated
- No proteinuria Not diagnosed with preeclampsia
(unaware of updated guidelines)
- Patient to left side Inappropriate BP assessment
- Reactive NST
- BP now 155/100
- Given Vicodin for HA ... better Ignored symptoms
- DC home

24 yo G1P0 at 34.1 weeks

- Presented to hospital with HA at 1400 (15 hrs. later)
- BP 175/105, 2+ protein
- NST NR
- Labs sent: plts=55K, Cr=1.6, AST/ALT=320/150, Fibrinogen=175, INR=1.4
- No BP meds
- Mag started , had seizure mid-dose
- C/S for fetal decelerations
- PP hemorrhage with DIC

24 yo G1P0 at 34.1 weeks

- Presented to hospital with HA at 1400
- BP 175/105, 2+ protein **Diagnosed with preeclampsia**
- NST NR
- Labs sent: plts=55K, Cr=1.6, AST/ALT=320/150, Fibrinogen=175, INR=1.4
- No BP meds **Diastolic BP < 110 but Systolic BP 175**
- Mag started , had seizure mid-dose **Too late**
- Emergent C/S for fetal decelerations **Indicated?**
- PP hemorrhage with DIC **Preventable**



Thank you

References

- Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstet Gynecol.* 2013 Nov;122(5):1122-1131. doi: 10.1097/01.AOG.0000437382.03963.88. PMID: 24150027.
- Committee on Obstetric Practice. Committee Opinion No. 692: Emergent Therapy for Acute-Onset, Severe Hypertension During Pregnancy and the Postpartum Period. *Obstet Gynecol.* 2017 Apr;129(4):e90-e95. doi: 10.1097/AOG.0000000000002019. PMID: 28333820.
- van Veen, Teelkien R., et al. "Cerebral autoregulation in normal pregnancy and preeclampsia." *Obstetrics & Gynecology* 122.5 (2013): 1064-1069.
- Council on Patient Safety in Women's Health Care Hypertension Patient Safety Bundle 2015
- High Risk & Critical Care Obstetrics 3rd ed. Troiano, Harvey, Flood- Chez
- Perinatal Nursing 4th ed. Simpson, Creehan, AWHONN
- Coppage, K, Sibai, B, Preeclampsia and Eclampsia. *Glob. libr. women's med.*, (ISSN: 1756-2228) 2008; DOI 10.3843/GLOWM.10158 Retrieved from Global Library of Women's Medicine. Updated 2017
- UpToDate: Preeclampsia Prevention. Feb. 2023
- Kilpatrick SJ, Abreo A, Greene N, Melsop K, Peterson N, Shields LE, Main EK. Severe maternal morbidity in a large cohort of women with acute severe intrapartum hypertension. *Am J Obstet Gynecol.* 2016 Jul;215(1):91.e1-7. doi: 10.1016/j.ajog.2016.01.176. Epub 2016 Jan 30. PMID: 26829504.
- Burton GJ, Redman CW, Roberts JM, Moffett A. Pre-eclampsia: pathophysiology and clinical implications. *BMJ.* 2019 Jul 15;366:l2381. doi: 10.1136/bmj.l2381. PMID: 31307997.
- Elgendy IY, Bukhari S, Barakat AF, Pepine CJ, Lindley KJ, Miller EC; American College of Cardiology Cardiovascular Disease in Women Committee. Maternal Stroke: A Call for Action. *Circulation.* 2021 Feb 16;143(7):727-738. doi: 10.1161/CIRCULATIONAHA.120.051460. Epub 2021 Feb 15. PMID: 33587666; PMCID: PMC8049095.